

PG No. 33 $D_{2d}(1)$ $\bar{4}2m$ (-4m2 setting) [tetragonal] (axial, internal polar dipole)

* Harmonics for rank 0

* Harmonics for rank 1

$$\bar{\mathbb{G}}_1^{(1,0)}[q](A_2)$$

** symmetry

$$z$$

** expression

$$\frac{\sqrt{2}Q_{xy}}{2} - \frac{\sqrt{2}Q_{yx}}{2}$$

$$\bar{\mathbb{G}}_{1,1}^{(1,0)}[q](E), \bar{\mathbb{G}}_{1,2}^{(1,0)}[q](E)$$

** symmetry

$$y$$

$$x$$

** expression

$$-\frac{\sqrt{2}Q_{xz}}{2} + \frac{\sqrt{2}Q_{zx}}{2}$$

$$\frac{\sqrt{2}Q_{yz}}{2} - \frac{\sqrt{2}Q_{zy}}{2}$$

* Harmonics for rank 2

$$\bar{\mathbb{G}}_2^{(1,0)}[q](A_1)$$

** symmetry

$$\sqrt{3}xy$$

** expression

$$-\frac{\sqrt{2}Q_{xxz}}{2} + \frac{\sqrt{2}Q_{yyz}}{2} + \frac{\sqrt{2}Q_z(x-y)(x+y)}{2}$$

$$\bar{\mathbb{G}}_2^{(1,0)}[q](A_2)$$

** symmetry

$$\frac{\sqrt{3}(x-y)(x+y)}{2}$$

** expression

$$\frac{\sqrt{2}Q_{xyz}}{2} + \frac{\sqrt{2}Q_{yxz}}{2} - \sqrt{2}Q_{zxy}$$

$$\bar{\mathbb{G}}_2^{(1,0)}[q](B_1)$$

** symmetry

$$-\frac{x^2}{2} - \frac{y^2}{2} + z^2$$

** expression

$$\frac{\sqrt{6}Q_{xyz}}{2} - \frac{\sqrt{6}Q_{yxz}}{2}$$

$$\bar{\mathbb{G}}_{2,1}^{(1,0)}[q](E), \bar{\mathbb{G}}_{2,2}^{(1,0)}[q](E)$$

** symmetry

$$\sqrt{3}yz$$

$$-\sqrt{3}xz$$

** expression

$$\frac{\sqrt{2}Q_x(y-z)(y+z)}{2} - \frac{\sqrt{2}Q_yxy}{2} + \frac{\sqrt{2}Q_zxz}{2}$$

$$-\frac{\sqrt{2}Q_xxy}{2} + \frac{\sqrt{2}Q_y(x-z)(x+z)}{2} + \frac{\sqrt{2}Q_zyz}{2}$$

* Harmonics for rank 3

$$\bar{\mathbb{G}}_3^{(1,0)}[q](A_2)$$

** symmetry

$$-\frac{z(3x^2 + 3y^2 - 2z^2)}{2}$$

** expression

$$-\frac{\sqrt{3}Q_{xy}(x^2 + y^2 - 4z^2)}{4} + \frac{\sqrt{3}Q_{yx}(x^2 + y^2 - 4z^2)}{4}$$

$$\vec{\mathbb{G}}_3^{(1,0)}[q](B_1)$$

** symmetry

$$\frac{\sqrt{15}z(x-y)(x+y)}{2}$$

** expression

$$\frac{\sqrt{5}Q_{xy}(x^2 - y^2 + 2z^2)}{4} - \frac{\sqrt{5}Q_{yx}(x^2 - y^2 - 2z^2)}{4} - \sqrt{5}Q_zxyz$$

$$\vec{\mathbb{G}}_3^{(1,0)}[q](B_2)$$

** symmetry

$$\sqrt{15}xyz$$

** expression

$$\frac{\sqrt{5}Q_{xx}(y-z)(y+z)}{2} - \frac{\sqrt{5}Q_{yy}(x-z)(x+z)}{2} + \frac{\sqrt{5}Q_{zz}(x-y)(x+y)}{2}$$

$$\vec{\mathbb{G}}_{3,1}^{(1,0)}[q](E, 1), \vec{\mathbb{G}}_{3,2}^{(1,0)}[q](E, 1)$$

** symmetry

$$-\frac{y(3x^2 - 2y^2 + 3z^2)}{2}$$

$$\frac{x(2x^2 - 3y^2 - 3z^2)}{2}$$

** expression

$$\frac{\sqrt{3}Q_{xz}(x^2 - 4y^2 + z^2)}{4} - \frac{\sqrt{3}Q_{zx}(x^2 - 4y^2 + z^2)}{4}$$

$$\frac{\sqrt{3}Q_{yz}(4x^2 - y^2 - z^2)}{4} - \frac{\sqrt{3}Q_{zy}(4x^2 - y^2 - z^2)}{4}$$

$$\vec{\mathbb{G}}_{3,1}^{(1,0)}[q](E, 2), \vec{\mathbb{G}}_{3,2}^{(1,0)}[q](E, 2)$$

** symmetry

$$\frac{\sqrt{15}y(x-z)(x+z)}{2}$$

$$\frac{\sqrt{15}x(y-z)(y+z)}{2}$$

** expression

$$-\frac{\sqrt{5}Q_{xz}(x^2 + 2y^2 - z^2)}{4} + \sqrt{5}Q_{yxyz} + \frac{\sqrt{5}Q_{zx}(x^2 - 2y^2 - z^2)}{4}$$

$$-\sqrt{5}Q_{xxyz} + \frac{\sqrt{5}Q_{yz}(2x^2 + y^2 - z^2)}{4} + \frac{\sqrt{5}Q_{zy}(2x^2 - y^2 + z^2)}{4}$$

* Harmonics for rank 4

$$\vec{\mathbb{G}}_4^{(1,0)}[q](A_1)$$

** symmetry

$$-\frac{\sqrt{5}xy(x^2 + y^2 - 6z^2)}{2}$$

** expression

$$\frac{Q_{xxz}(x^2 + 15y^2 - 6z^2)}{4} - \frac{Q_{yyz}(15x^2 + y^2 - 6z^2)}{4} - \frac{Q_z(x-y)(x+y)(x^2 + y^2 - 6z^2)}{4}$$

$$\vec{\mathbb{G}}_4^{(1,0)}[q](A_2)$$

** symmetry

$$\frac{\sqrt{5}(x-y)(x+y)(x^2+y^2-6z^2)}{4}$$

** expression

$$-\frac{Q_x y z (3x^2 - 4y^2 + 3z^2)}{2} + \frac{Q_y x z (4x^2 - 3y^2 - 3z^2)}{2} - \frac{Q_z x y (x^2 + y^2 - 6z^2)}{2}$$

$$\vec{\mathbb{G}}_4^{(1,0)}[q](B_1, 1)$$

** symmetry

$$\frac{\sqrt{21}(x^4 - 3x^2 y^2 - 3x^2 z^2 + y^4 - 3y^2 z^2 + z^4)}{6}$$

** expression

$$-\frac{\sqrt{105}Q_x y z (y-z)(y+z)}{6} + \frac{\sqrt{105}Q_y x z (x-z)(x+z)}{6} - \frac{\sqrt{105}Q_z x y (x-y)(x+y)}{6}$$

$$\vec{\mathbb{G}}_4^{(1,0)}[q](B_1, 2)$$

** symmetry

$$-\frac{\sqrt{15}(x^4 - 12x^2 y^2 + 6x^2 z^2 + y^4 + 6y^2 z^2 - 2z^4)}{12}$$

** expression

$$-\frac{\sqrt{3}Q_x y z (9x^2 + 2y^2 - 5z^2)}{6} + \frac{\sqrt{3}Q_y x z (2x^2 + 9y^2 - 5z^2)}{6} + \frac{7\sqrt{3}Q_z x y (x-y)(x+y)}{6}$$

$$\vec{\mathbb{G}}_4^{(1,0)}[q](B_2)$$

** symmetry

$$\frac{\sqrt{35}x y (x-y)(x+y)}{2}$$

** expression

$$-\frac{\sqrt{7}Q_x x z (x^2 - 3y^2)}{4} + \frac{\sqrt{7}Q_y y z (3x^2 - y^2)}{4} + \frac{\sqrt{7}Q_z (x^2 - 2xy - y^2)(x^2 + 2xy - y^2)}{4}$$

$$\vec{\mathbb{G}}_{4,1}^{(1,0)}[q](E, 1), \vec{\mathbb{G}}_{4,2}^{(1,0)}[q](E, 1)$$

** symmetry

$$\frac{\sqrt{35}y z (y-z)(y+z)}{2}$$

$$-\frac{\sqrt{35}x z (x-z)(x+z)}{2}$$

** expression

$$\frac{\sqrt{7}Q_x (y^2 - 2yz - z^2)(y^2 + 2yz - z^2)}{4} - \frac{\sqrt{7}Q_y x y (y^2 - 3z^2)}{4} + \frac{\sqrt{7}Q_z x z (3y^2 - z^2)}{4}$$

$$-\frac{\sqrt{7}Q_x x y (x^2 - 3z^2)}{4} + \frac{\sqrt{7}Q_y (x^2 - 2xz - z^2)(x^2 + 2xz - z^2)}{4} + \frac{\sqrt{7}Q_z y z (3x^2 - z^2)}{4}$$

$$\vec{\mathbb{G}}_{4,1}^{(1,0)}[q](E, 2), \vec{\mathbb{G}}_{4,2}^{(1,0)}[q](E, 2)$$

** symmetry

$$\frac{\sqrt{5}y z (6x^2 - y^2 - z^2)}{2}$$

$$\frac{\sqrt{5}x z (x^2 - 6y^2 + z^2)}{2}$$

** expression

$$\frac{Q_x (y-z)(y+z)(6x^2 - y^2 - z^2)}{4} - \frac{Q_y x y (6x^2 - y^2 - 15z^2)}{4} + \frac{Q_z x z (6x^2 - 15y^2 - z^2)}{4}$$

$$\frac{Q_x x y (x^2 - 6y^2 + 15z^2)}{4} - \frac{Q_y (x-z)(x+z)(x^2 - 6y^2 + z^2)}{4} - \frac{Q_z y z (15x^2 - 6y^2 + z^2)}{4}$$